



## SPECC1L gene

sperm antigen with calponin homology and coiled-coil domains 1 like

### Normal Function

The *SPECC1L* gene provides instructions for making a protein called cytospin-A. This protein stabilizes components of the cell's structural framework (cytoskeleton) called microtubules, which are rigid, hollow fibers that help maintain the cell's shape. Stabilization of microtubules is necessary for these fibers to regulate various cell processes including the movement of cells to their proper location (cell migration). In order for cells to move, microtubules elongate in a specific direction, changing the shape of the cytoskeleton and allowing the cell to move in that direction. Migration of cells to their proper location during development ensures normal tissue formation.

During development of the embryo, cytospin-A plays a role in the migration of cells called neural crest cells, which originate in the developing spinal cord and migrate to specific regions in the embryo to form different structures. Cytospin-A is specifically involved in the migration of neural crest cells that come together to form the forehead, nasal bridge, and lower jaw.

### Health Conditions Related to Genetic Changes

#### Opitz G/BBB syndrome

At least two mutations in the *SPECC1L* gene have been found to cause Opitz G/BBB syndrome. This condition causes several abnormalities along the midline of the body, including widely spaced eyes (ocular hypertelorism), throat malformations that can cause difficulty breathing or swallowing, brain malformations, distinct facial features, and genital abnormalities in males. The *SPECC1L* gene mutations that cause Opitz G/BBB syndrome change single protein building blocks (amino acids) in the cytospin-A protein, reducing the protein's ability to interact with components of the cytoskeleton. As a result, microtubules are disorganized and cells have trouble migrating to their proper location. Because the *SPECC1L* gene plays a role in facial development, mutations in this gene likely account for an opening in the lip (cleft lip) with or without an opening in the roof of the mouth (cleft palate) seen in some individuals with Opitz G/BBB syndrome. However, it is unclear how *SPECC1L* gene mutations cause the other features of this disorder.

#### other disorders

Mutations in the *SPECC1L* gene have also been found to cause openings in facial structures called oblique facial clefts, also known as Tessier clefts. These clefts are


similar to cleft lip and palate, but are more severe. The facial clefts can occur on one or both sides of the face; they typically involve the mouth and can include tissues up to the eyes. Health problems experienced by affected individuals depend on the severity of the cleft, but generally include problems eating and breathing.

The *SPECC1L* gene mutations that cause oblique facial clefts diminish the function of the cytospin-A protein. As a result, the protein cannot interact with the cytoskeleton, and microtubules are disorganized. These problems with the cytoskeleton impair cell migration, particularly of cells that come together to form the structures of the face. A decrease in cytospin-A function impairs cell migration during facial development, leaving openings, or clefts, where tissue should normally be.

### Chromosomal Location

Cytogenetic Location: 22q11.23, which is the long (q) arm of chromosome 22 at position 11.23

Molecular Location: base pairs 24,270,817 to 24,417,740 on chromosome 22 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

### Other Names for This Gene

- cytokinesis and spindle organization A
- cytospin A
- CYTSA
- KIAA0376
- sperm antigen with calponin homology and coiled-coil domains 1-like

### Additional Information & Resources

#### Educational Resources

- The Cell: A Molecular Approach (second edition, 2000): Microtubules  
<https://www.ncbi.nlm.nih.gov/books/NBK9932/>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28SPECC1L%5BTIAB%5D%29+AND+english%5Bla%5D>

### OMIM

- FACIAL CLEFTING, OBLIQUE, 1  
<http://omim.org/entry/600251>
- SPERM ANTIGEN WITH CALPONIN HOMOLOGY AND COILED-COIL DOMAINS 1-LIKE  
<http://omim.org/entry/614140>

### Research Resources

- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=SPECC1L%5Bgene%5D>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=29022](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=29022)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/23384>
- UniProt  
<http://www.uniprot.org/uniprot/Q69YQ0>

### **Sources for This Summary**

- Gfrerer L, Shubinets V, Hoyos T, Kong Y, Nguyen C, Pietschmann P, Morton CC, Maas RL, Liao EC. Functional analysis of SPECC1L in craniofacial development and oblique facial cleft pathogenesis. *Plast Reconstr Surg*. 2014 Oct;134(4):748-59. doi: 10.1097/PRS.0000000000000517.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/25357034>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4430087/>
- Kruszka P, Li D, Harr MH, Wilson NR, Swarr D, McCormick EM, Chiavacci RM, Li M, Martinez AF, Hart RA, McDonald-McGinn DM, Deardorff MA, Falk MJ, Allanson JE, Hudson C, Johnson JP, Saadi I, Hakonarson H, Muenke M, Zackai EH. Mutations in SPECC1L, encoding sperm antigen with calponin homology and coiled-coil domains 1-like, are found in some cases of autosomal dominant Opitz G/BBB syndrome. *J Med Genet*. 2015 Feb;52(2):104-10. doi: 10.1136/jmedgenet-2014-102677. Epub 2014 Nov 20.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/25412741>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4393015/>

- OMIM: SPERM ANTIGEN WITH CALPONIN HOMOLOGY AND COILED-COIL DOMAINS 1-LIKE  
<http://omim.org/entry/614140>
  - Saadi I, Alkuraya FS, Gisselbrecht SS, Goessling W, Cavallero R, Turbe-Doan A, Petrin AL, Harris J, Siddiqui U, Grix AW Jr, Hove HD, Leboulch P, Glover TW, Morton CC, Richieri-Costa A, Murray JC, Erickson RP, Maas RL. Deficiency of the cytoskeletal protein SPECC1L leads to oblique facial clefting. *Am J Hum Genet.* 2011 Jul 15;89(1):44-55. doi: 10.1016/j.ajhg.2011.05.023. Epub 2011 Jun 23.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/21703590>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3135813/>
- 

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/SPECC1L>

Reviewed: January 2015  
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services